

Journal of The International Association of Advanced Technology and Science**INVERTEBRATE LECTINS AND THEIR BIOMEDICAL APPLICATIONS****Prakash Shoba¹. S, Basil Rose². M.R and Delphine Rose. M.R¹**

¹Department of Zoology, Jayaraj Annapackiam College for Women, Periyakulam, Theni
prakash.shoba061@gmail.com

²Department of Zoology, Holy Cross College, Nagercoil

ABSTRACT

Invertebrate lectins are diverse biological functions, including “self-nonsel” recognition, bacterial agglutination and lysis, metamorphosis, moulting, wound repair, regeneration and host parasite interactions. Agglutinins/lectins are predominantly found in the hemolymph, hemocytes and hepatopancreas of crustaceans. The functioning of the agglutinins/lectins depends upon its ability to recognize and bind to specific sugars on the cell surface glyco-conjugates. Lectins can be used as tools for blood typing, diagnosing of micro-organisms, mitogenic stimulation of lymphocytes, and discrimination between normal and malignant cells, purification of glyco-conjugates and as tools to examine cell surface carbohydrates. The functioning of the agglutinin depends upon its ability to recognize and bind to specific sugars on the cell surface glyco-conjugates.

KEYWORDS

Lectins, agglutination, glyco-conjugates, Invertebrates, tumor.

INTRODUCTION

The capacity to mount an immune response that eliminates infection of a host by a microbial pathogen is critical for species survival and propagation¹. Invertebrate animals, which lack adaptive immune systems, have developed other systems of biological host defense, so called innate immunity, that respond to common antigens on the cell surfaces of potential pathogens. The innate immune system is the first line of inducible host defense against bacterial, fungal and viral pathogens². This defense system is essential for the survival and perpetuation of all multicellular organisms^{3,4}. Invertebrates which do not possess immunoglobulins, have developed unique modalities, to detect and respond to microbial surface antigens like lipopolysaccharides (LPS), lipoteichoic acids, lipoproteins, peptidoglycans (PGN) and (1-3) β -D-glucans⁵.

Defense molecules in invertebrates

Invertebrate innate immunity relies on both cellular and humoral components. Invertebrate humoral immunity involves the presence of biologically active molecules that occur naturally or that may be induced. These molecules, by their lytic or agglutinating properties are able to act on the antigens responsible for their induction; only in this respect do they resemble vertebrate antibodies. The humoral factors (native, induced) in invertebrates include LPS binding proteins, phenoloxidase system, antibacterial proteins, antifungal proteins, lysins and agglutinins. According to Iwanaga and Lee (2005)⁶, the defense molecules include phenoloxidases, clotting factors, complement factors, protease inhibitors, antimicrobial peptides, toll-free receptors, lectins and other humoral factors found mainly in hemolymph plasma and cell hemocytes. Hemolymph in association with the hemocytes performs an essential role in immune defense and wound healing in insects and other invertebrates, revealing that both cellular (phagocytosis, nodule formation, encapsulation) and humoral elements (agglutinin, antibacterial proteins) are involved in the reaction to non-self⁷.

Agglutinins/Lectins

Lectin from castor bean extracts were first described by Stillmark (1888)⁸. They are distributed widely in bacteria, plant and animals⁹ and have been given various definitions. Lectins

are a group of proteins that interact with glycoproteins and glycolipids by binding to specific carbohydrate residues¹⁰. Lectins (formerly found as hemagglutinin), are carbohydrate-binding proteins which recognize specific carbohydrate structures on mammalian cells¹¹. Lectins are a group of sugar binding proteins that recognize a specific carbohydrate structure and agglutinate various cells by binding to cell surface glyco-conjugates¹². Lectins are protein complexes with carbohydrate specific binding properties that have been found in viruses, bacteria, plants, invertebrates and vertebrates and may have a wide variety of functions¹³.

Sialic acids

Sialic acids are a family of nine carbon acidic ketoses found predominantly at the non reducing end of oligosaccharide chains on glycoproteins and glycolipids¹⁴. Sialic acids show remarkable structural diversity with the family currently comprising over 50 naturally occurring members¹⁵. The largest structural variations of naturally occurring sialic acids are at carbon 5, which can be substituted with an acetamino, hydroxyl acetamino or hydroxyl moiety to form 5-N-acetyl neuraminic acid (Neu5Ac), 5-N-glycolyl neuraminic acid (Neu5Gc) or Keto Deamino Neuraminic acid (KDN) respectively¹⁶. This diversity of structure reflects their involvement in a variety of biologically and medically important functions¹⁷.

Importance of sialic acids

Sialic acids play an important role as ligands in cell sociology. The unique structural features of the molecule, which includes a negative charge owing to a carboxyl group, enables it to play a role in cellular functions, such as transport of positively charged compounds, cell to cell repulsion, influencing conformation of glycoprotein on cell membranes, and even masking antigenic determinants on receptor molecules¹⁸. The derivatives of sialic acids are very important constituents of the cell surface. The normal human tissues possess the common sialic acid, N-Acetyl neuraminic acid, which gets modified to O-Ac or NeuGc on neoplastic transformation¹⁹. They occur as components of glycoproteins and glycosphingolipids on the cell surface and in body fluids. Sialylation of glycoprotein changes under pathological

conditions as well as during developmental stages and altered sialylation often has significant implications in the physiological role of glycoproteins²⁰. Lectins that recognize the linkages or modifications of sialic acids are therefore indispensable as reagents in biochemical research and diagnostic analysis.

Mode of action of lectins

Invertebrate agglutinins are sugar-binding proteins with multiple binding sites, diverse biological roles²¹ and biomedical applications. They may recognize a part of a sugar²², a whole sugar²³, their glycosidic linkage²⁴ or a sequence of sugars²⁵. Lectins, by acting as opsonins may mediate phagocytosis of foreign particles by hemocytes²⁶. In invertebrates, which do not have an antibody based immune system²⁷, lectin may act as recognition molecules for defense activities such as leukocyte aggregation, heteroagglutination and opsonization²⁸. Lectins play crucial roles in innate immunity and host defense both in vertebrates and invertebrates with involvement in processes such as non-self recognition, inflammation, opsonization, cell-cell or cell-extra cellular matrix interaction, fertilization, development and regeneration, cell aggregation, wound repair, metamorphosis and transport of complex sugars²⁹.

Biomedical application of Lectins/Agglutinins

The availability of a number of lectins with diverse carbohydrate specificities makes these proteins a useful tool in biochemistry, immunology and cellular biology. Lectins are very sensitive to small variations in tissue epitopes and the affinity of the lectin for its binding sugars in a tissue is also affected by surrounding carbohydrate. The sialic acid binding specificity of this lectin was used to separate immature mouse thymocytes (low sialic acid content) from mature thymocytes (high sialic acid content)³⁰.

Typical recognition molecules, the mannose binding lectin (MBL) and ficolins, are involved in the specific recognition of carbohydrates on pathogenic micro-organism including bacteria, fungi, parasitic protozoans and viruses³¹. Ficolins and collectins play important role in host defense, non-self recognition and opsonic activity of the neutrophil³². Lectins are used to identify

inflammation, cancer, pregnancy and foetal development³³. Tumor immunologists have developed murine and human monoclonal antibodies against sialyl antigens for diagnosis and immunotherapy³⁴. Lectin affinity chromatography provides evidence for the alteration in the normal glycosylation of the alpha subunit of the human chorionic gonadotropic hormone in patients with pituitary tumors³⁵.

Lectins recognize sugars that constitute the surface components of pathogenic bacteria that invade the body cavity³⁶. With a battery of immobilized lectins, mixtures of glycopeptides or oligosaccharides obtained by enzymatic or chemical cleavage of purified glycoproteins can be separated into homogenous compounds³⁷. Lectins are used to characterize the structural changes on oligosaccharides in leukemias. The lectin from *M. rosenbergii* could be considered a useful tool for the diagnosis and study of T-cell acute lymphoblastic leukemia³⁸.

Several sialic acid specific lectins purified from plant and other sources have been employed as tools for the detection and separation of sialic-acid containing glyco-conjugates³⁹. Lectins are used in cell separation, mitogenic stimulation and blood grouping⁴⁰. Lectins could be used as drug carriers for carrying biologically active molecules and to direct those to specific molecules and cell organelles⁴¹.

Lectins have been implicated in the attachment of a sperm to an egg, the first step in fertilization⁴². Lectins are used to study changes in the distribution of glyco-conjugates during embryonic development⁴³. Forensic science laboratories use immobilized lectins for rapid typing of haptoglobin in blood stains⁴⁴. Lysosomal storage diseases characterized by accumulation of corresponding substances could be revealed by lectin staining⁴⁵. Lectin is used as a tracer for mapping neuronal connections in neuroanatomy⁴⁶. Cells pre-treated with anti-lectin antibody showed decreased metastatic potential⁴⁷. Protein-carbohydrate interaction plays a role in identifying the spread of cancer cells from the main tumor throughout the body⁴⁸.

Lectins present on various human and murine metastatic tumor cells influence the pathogenesis of cancer metastasis⁴⁹. Lectins are specific in their reactions with human blood groups (ABO and MN) and sub groups (A₁) and have therefore been used in blood typing and in investigation of the chemical basis of blood group specificity⁵⁰. The stimulation of lymphocytes by lectin also provide an important tool for the examination of the biochemical events involved in the conversion of a resting cell into an actively growing one⁵¹. Lectins (*Macrobrachium* lectin) are used in studying the process of protein glycolysation in the brain from patients with Alzheimer's disease⁵².

N-acetyl-D-glucosamine specific lectin in crustacean hemocytes is used for wound healing, pathogen encapsulation, and maintenance of normal crustacean connective tissues⁵³. A novel lectin (EIL) isolated from the marine hair crab *Erimuramus isenbeckii*, with high specificities for both O-acetyl sialic acid and mannose that are present in bacterial pathogens suggests that EIL may act as a defense protein against infection in this crab⁵⁴. Romano and Alberto (2004)⁵⁵ have showed that the incorporation of the purified lectin from *Macrobrachium rosenbergii* in the food composition induced resistance of the specimen, *Litopenaeus vannamei* against the white spot disease virus.

Conclusion

Although a number of studies have demonstrated the presence of humoral agglutinins in several crustacean species, it can be noted that the immunological role of these lectins/agglutinins remain unknown and that the carbohydrate specificity of serum agglutinin from crustaceans have been elucidated only in a few species.

Acknowledgement

I greatly acknowledge my friends A. Punitha and S. Mary Mettilda Bai for their support towards my research work

Reference

1. Young, J.A.T. and Dillin, A. (2004). Mapping innate immunity. *Proceedings of the National academy of Sciences, USA*. 101: 12781-12782.
2. Hoebe, K., Jansen, E. and Beutler, B. (2004). The interface between innate and adaptive immunity. *Nat. Immunol.* 5: 971-974.
3. Hoffmann, J.A., Kafatos, F.C. Janeway, C.A. Jr. and Ezekowitz, R.A.B. (1999). Phylogenetic perspectives in innate immunity. *Science*, 284: 1313-1318.
4. Salzet, M. (2001). Vertebrate innate immunity resembles a mosaic of invertebrate immune responses. *Trends Immunol.* 22: 285-288.
5. Begum, N., Matsumoto M., Tsuji, S., Toyoshima, K. and Seya, T. (2000). The primary host defense system across humans, flies and plants. *Current Trends in Immunology*, 3: 59-74.
6. Iwanaga, S. and Lee, B.L. (2005). Recent advances in the innate immunity of invertebrates animals. *J. Biochem. Mol. Biol.* 38(2): 128-150.
7. Gupta, A.P. (1986). *Hemocytic and Humoral Immunity in Arthropods*. J. Wiley, New York.
8. Stillmark, H. (1888). *Über rizin ein giftiges ferment aus samen von Ricinus communis L., und ainigen Euphorbiaceen*. Dorpat (Tartu). Inaugural dissertation.
9. Liener, I.E., Sharon, N. and Goldstein, I.J. (1986). In: "The Lectins: Properties, Functions and Applications in Biology and medicine". Academic Press, NY. pp. 11-15.
10. Goldstein, I.J. and Hayes, C.E. (1978). The lectins: carbohydrate binding of plants and animals. *Adv. Carbohydr. Chem. Biochem.* 35: 127-340.
11. Nangia-Makker, P., Honjo, Y., Sarvis, R., Akahani., Hogan, V. and Pienta, K. J. (2000). Galactin-3 induces endothelial cell morphogenesis and angiogenesis. *Am. J. Pathol.* 156: 899-909.
12. Matsubara, H., Nakamura, S., Hirabayashi, J., Jimbo, M., Kamiya, H., Ogawa, T. and Muramoto, K. (2007). Diverse sugar-binding specificities of marine invertebrate C-Type lectins. *Biosci. Biotechnol. Biochem.* 71(2): 513-519.
13. Sharon, N. and Lis, H. (1972). Cell-agglutinating and sugar-specific proteins. *Science*, 177 (4053): 949-959.
14. Lehman, F., Tiralongo, E and Tiralongo, J. (2006). Sialic acid specific lectins: occurrence, specificity and function. *Cell. Mol. Life Sci.* 63: 1331-1354.
15. Angata, T and Varki, A. (2002). Chemical diversity in the sialic acids and related alpha-keto acids: an evolutionary perspective. *Chem. Rev.* 102: 439-469. (Schauer and Kamerling, 1997)

17. Jeanloz, R.W. and Cudington, J.F. (1976). The biological role of sialic acid at the surface of the cell. In *Biological Roles of Sialic Acid* (Rosenberg A, Schenck (Eds), Plenum Press, New York, 201-238.
18. Narayanas, S. (1994). Sialic acid as a tumor marker. *Animals of Clin. And Lab. Science*, 24(4): 376-384.
19. Ravindranath, M.H. and Irie, R.F. (1988). Gangliosides as antigens of human melanoma in "Malignant melanoma: biology, diagnosis and therapy" (Nathanson, Lad) Kluwer Acad, Puplishers Boston, pp. 17-43.
20. Varki, A. (1997). Sialic acids as ligands in recognition phenomena. *FASEB J*. 248-255.
21. Kabiler, D. and Mirelman, D. (1980). Lectin activity in *Entamoeba histolytica* trophozoites. *Infect. Immun.* 29: 221-225
22. Ravindranath, M.H., Higa, H.H., Cooper, E.L. and Paulson, J.C. (1985). Purification and characterization of an O-acetyl sialic acid specific lectin from the crab *Cancer antennarius*. *J. Biol. Chem.* 260: 8850-8856
23. Bretting, H. and Kabat, E.A. (1976). Purification and characterization of the agglutinin from the sponge *Asimella polyopoides* and a study of their combining sites. *Biochemistry*, 15: 3228- 3236.
24. Koch, O.M., Lee, C.K. and Uhlenbruck, G. (1982). Cerianthin lectins: A new group of agglutinins from *Cerianthus membranaceus* (Singapore). *Immunol. Biol.* 163: 53-62.
25. Kabiler, D. and Mirelman, D. (1980). Lectin activity in *Entamoeba histolytica* trophozoites. *Infect. Immun.* 29: 221-225
26. Vasta, R.W. (1981). Invertebrate lectins. II. Diversity of specificity, biological synthesis and function in recognition. *Dev. Comp. Immunol.* 5: 536-545.
27. Warr, G.W. (1981). Immunity in invertebrates. *J. Invertebr. Pathol.* 38: 311-314.
28. Vasta, G.R. and Marchalonis, J.J. (1984). Immunological significance of invertebrate lectins. *J. Comp. Physiol. B* 160: 119-126.
29. Ravindranath, M.H., Higa, H.H., Cooper, E.L. and Paulson, J.C. (1985). Purification and characterization of an O-acetyl sialic acid specific lectin from the crab *Cancer antennarius*. *J. Biol. Chem.* 260: 8850-8856.
30. Abel, C.A., Campbell, P.A., Vander Wall, J. and Hartman, A.L. (1984). Studies on the structure and carbohydrate binding properties of lobster agglutinin 1 (LAG-1), a sialic acid-binding lectin. *Prog. Clin. Biol. Res.* 157: 103-114.
31. Holmster, V., Thiel, S. and Jensenius, J.C. (2003). Collectins and Ficolins: Humoral lectins of the innate immune defense. *Annu. Rev. Immunol.* 21: 547-578.
32. Matsuhashi, M. (1996). The lectin pathway of the complement system. *Microbiol. Immunol.* 40: 887-893

33. Lampreave, F., Alva, M. and Pineiro, A. (1993).Cancavalin A crossed affinoelectrophoretic analysis of the major pig serum proteins during foetal development.Electrophoresis. 14 (3): 2149.
34. Ravindranath, M.H. and Irie, R.F. (1988). Gangliosides as antigens of human melanoma in "Malignant melanoma: biology, diagnosis and therapy".(Natharson, Led) Kluwer Acad, Puplishers Boston. pp. 17-43,
35. Endo, T., Nishimura, R., and Muchizuki, M. (1988).Altered glycosylation is induced in both α and β - subunits of chorionic gonadotropin produced by choricarcinoma. J. Biochem. (Japan) 103: 1035- 38.
36. Ravindranath, M.H. and Cooper, E.L. (1984).Crab lectins: receptor specificity and biomedical applications.Prog. Clin. Biol. Res. 157: 83-96.
37. Osawa, T. and Tsuji, T. (1987).Fractionation and structural assessment of oligosaccharides and glycopeptides by the use of immobilized lectin.Ann. Rev. Biochem. 56: 21-42.
38. Perez-Campos-Mayoral, L., Ruiz-Arguelles, A., Perez-Romano, B., Zenteno, E., Hernandez-Cruz, P., Marti Cruz, R. Martinz-Cruz, M., Pina-Canseco, S. ana Perez-Compos, E. (2008).Potential use of the *Macrobrachium rosenbergii* lectin for diagnosis of T-cell acute lymphoblastic leukemia.Tohoku. J. Exp. Med. 214(1): 11-16.
39. O'Reilly, D.R., Miller, L.K. and Lucknow, V.A. (1994).Baculovirus Expression Vectors: A Laboratory Manual, Oxford University Press, New York.
40. Jacobson, R.L. and Doyle, R.J. (1996). Lectin-parasite intractions.Parasitol. Today, 12:55-60.
41. Lavelle, E.C. (2001).Targeted delivary of drugs to the gastro intestinal tract.Crit. Rev. Ther. Drug Carrier Syst. 18 (4): 341-386.
42. Wassarman, P.M. (1987).The biology and chemistry of fertilization.Science, 235: 553-60.
43. D' Amico, P. and Jacobs, J.R. (1995). Lectin histochemistry of the *Drosophila* embryos.Tissue Cell, 27 (1): 23- 30.
44. Harada, A., Umetsu, Ikeda, N. and Suzuki, T. (1987).A new method for typing haptoglobin in blood stains using immobilized allo A lectin.J. Forensic Sci. 32(4): 1062- 1064.
45. Castagnaro, M., Alroy, J., Ucci, A.A. and Jaffe, R. (1987).Lectin histochemistry and Ultra structure of kidneys for patients with I- cell disease. Arch. Pathol. Lab. Med., 111: 285-90 46. (Sawchenko and Gerfen
47. Raz, A. and Lotan, R. (1987).Endogenous galactoside-binding lectins : a new class of functional tumor surface molecules related to metatodis. Cancer Metast. Rev. 6: 4333-52.

48. Shirahama, T., Ikoma, M., Muramatsu, T. and Ohi, Y. (1992). Expression of SSEA-1 carbohydrate antigen correlates with stage grade and metastatic potential of transitional cell carcinoma of the bladder. *J. Urol.* 148(4): 1319-1322.
49. Keida, and Monsigny, M. (1986). Involvement of membrane sugar receptor and membrane glycoconjugate in the adhesion of 3LL cell subpopulation to cultures pulmonary cells. *Invas. Metast.* 6: 347- 66.
50. Boyd, W.C. (1963). The lectins: their present status. *Vox Sang.* 8: 1-32.
51. Nowell, P.C. (1960). Phytohemagglutinin an initiator of mitosis in culture of animal and human leukocytes. *Cancer Res.* 20: 462-466.
52. Espinosa, B., Zenteno, R., Mena, R., Robitaille, Y., Zenteno, E. and Guevara, J. (2001). O-glycosylation in sprouting neurons in Alzheimer disease, indicating reactive plasticity. *J. Neuropathol. Exp. Neurol.* 60(5): 441-448.
53. Martin, G.G., Castro, C., Moy, N. and Rubin, N. (2000). N-acetyl-D-glucosamine in crustacean hemocytes; possible functions and usefulness in hemocyte classification. *J. Invertebr. Biol.* 122(3): 265-270.
54. Na, YJ., Kim, YJ., Park, BT., Junk, BW., Hwang, KW. and Kim, H. (2007). A novel lectin isolated from the hemolymph of the marine hair crab *Erimacrus isenbeckii*. *Protein Pept Lett.* 14(8): 800-803.
55. Romano and Alberto, L. (2004). Substantially pure lectins for prevention and treatment of crustacean infected by baculovirus, a food composition containing the lectin and a method for increasing the crustacean resistance to infections. United States Patent 20040102363.